arXiv:1005.3926v2 [math.CO] 24 Aug 2010

# On the Multi-coloured Ramsey Numbers of Cycles

Tomasz Łuczak [*][†] Miklós Simonovits [‡][§] Jozef Skokan [¶]

August 21, 2010

**Abstract**

For a graph $L$ and an integer $k \geq 2$, $R_k(L)$ denotes the smallest integer $N$ for which for any edge-colouring of the complete graph $K_N$ by $k$ colours there exists a colour $i$ for which the corresponding colour class contains $L$ as a subgraph.

Bondy and Erdős conjectured that, for an odd cycle $C_n$ on $n$ vertices,

$$R_k(C_n) = 2^{k-1}(n-1) + 1 \quad \text{for } n > 3.$$

They proved the case when $k = 2$ and also provided an upper bound $R_k(C_n) \leq (k+2)!n$. Recently, this conjecture has been verified for $k = 3$ if $n$ is large. In this note, we prove that for every integer $k \geq 4$,

$$R_k(C_n) \leq k2^k n + o(n), \qquad \text{as } n \to \infty.$$

When $n$ is even, Yongqi, Yuansheng, Feng, and Bingxi gave a construction, showing that $R_k(C_n) \geq (k-1)n - 2k + 4$. Here we prove that if $n$ is even, then

$$R_k(C_n) \leq kn + o(n), \qquad \text{as } n \to \infty.$$

---

[*] Faculty of Mathematics and Computer Science, Adam Mickiewicz University, 61-614 Poznań, Poland, e-mail: `tomasz@amu.edu.pl`

[†] The author was partially supported by the Foundation for Polish Science.

[‡] Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences H-1053 Budapest, Reáltanoda u. 13-15., Hungary, e-mail: `miki@renyi.hu`

[§] The author was partially supported by the Hungarian National Science Foundation grants OTKA T 026069, T 038210, T 0234702, and T 69062.

[¶] Department of Mathematics, London School of Economics, Houghton Street, London WC2A 2AE, United Kingdom, e-mail: `jozef@member.ams.org`

# 1. Introduction

In this note we shall consider Ramsey problems connected to edge-colourings of ordinary graphs with $k$ colours, for a given $k \geq 2$, and try to ensure monochromatic cycles of a given length. We shall use the standard notation. Given a graph $G = (V, E)$, $v(G)$ denotes the number of vertices and $e(G)$ the number of edges in $G$. For a subset $W$ of $V$, $G[W]$ is the subgraph of $G$ induced by the vertices in $W$.

For graphs $L_1, \ldots, L_k$, the Ramsey number $R(L_1, \ldots, L_k)$ is the minimum integer $N$ such that for any edge-colouring of the complete graph $K_N$ by $k$ colours there exists a colour $i$ for which the $i^{\text{th}}$ colour class contains $L_i$ as a subgraph. For $L_1 = L_2 = \ldots = L_k = L$, we set $R_k(L) := R(L_1, \ldots, L_k)$.

The behaviour of the Ramsey number $R(C_n, C_m)$ has been studied by several authors, for example, by Bondy and Erdős, [3], Faudree and Schelp, [6], Rosta, [13], and it is completely described and well-understood. Among others, it is known that

$$R_2(C_n) = \begin{cases} 2n - 1, & \text{if } n \geq 5 \text{ is odd,} \\ \frac{3n}{2} - 1, & \text{if } n \geq 6 \text{ is even.} \end{cases}$$

Bondy and Erdős [3] conjectured that $R_k(C_n) = 2^{k-1}(n-1)+1$ for every odd $n > 3$. The conjectured extremal colouring, giving the lower bound, can be easily constructed recursively: for two colours, take two disjoint sets of size $n-1$, colour all the pairs within each set by colour 1, and colour all the pairs joining these two sets by colour 2. For $i = 3, \ldots, k$, take two disjoint copies of the colouring for $i - 1$ colours and colour all the pairs joining these two copies by colour $i$. The final $k$-colouring has $2^{k-1}(n - 1)$ vertices and every monochromatic component has either only $n - 1$ vertices or it is bipartite and, therefore, does not contain odd cycles.

As for the upper bound for $R_k(C_n)$, $k \geq 3$, Łuczak [12] proved that if $n$ is odd, then $R_3(C_n) = 4n + o(n)$, as $n \to \infty$. Later, Kohayakawa, Simonovits, and Skokan [8, 9] showed that $R_3(C_n) = 4n - 3$ for all odd, sufficiently large values of $n$. The conjecture is still open for $k \geq 4$. Bondy and Erdős [3] remarked that they could prove $R_k(C_n) \leq (k + 2)!n$ for $n$ odd. In this note we shall give an upper bound which is correct up to $O(k)$ factor.

**Theorem 1.** *For every $k \geq 4$ and odd $n$,*

$$R_k(C_n) \leq k2^k n + o(n), \quad \text{as } n \to \infty.$$

The Ramsey number $R_k(C_n)$ behaves rather differently for even values of $n$. From [6] and [13], we know that $R_2(C_n) = 3n/2-1$ and, for large even $n$, Benevides and Skokan [2] proved that $R_3(C_n) = 2n$. Yongqi, Yuansheng, Feng, and Bingxi [15] gave a construction yielding

$$R_k(C_n) \geq (k-1)n - 2k + 4.$$

Here we prove the following.

**Theorem 2.** *For every $k \geq 2$ and even $n$,*

$$R_k(C_n) \leq kn + o(n), \qquad \text{as } n \to \infty.$$

In both results above, $k$ is fixed and $n$ is large. When $n$ is fixed and $k$ is large, we know even less. For $n$ odd and $k$ large, the original Bondy-Erdős bound from [3]

$$2^{k-1}(n-1) + 1 \leq R_k(C_n) \leq (k+2)!n$$

is still the best bound we have. There are better bounds for small values of $n$. For $n = 3$, it is known that

$$c_1 1073^{k/6} \leq R_k(C_3) \leq c_2 k!$$

with $c_2 < e$, see [1, 4, 16, 17]. For $n = 5$, Li [10] proved that

$$R_k(C_5) \leq \sqrt{18^k k!}, \qquad \text{for } k \geq 2.$$

For a fixed even $n$, it is known that

$$R_k(C_n) \leq ck^{\frac{n}{n-2}},$$

where $c$ is a constant depending only on $n$. Li and Lih [11] showed that this bound is asymptotically correct for $n = 4, 6, 10$.

## 2. Tools

We shall make use of the following result of Erdős and Gallai, [5].

**Theorem 3.** *Let $n \geq 3$. For any graph $G$ with at least $(n-1)(v(G)-1)/2+1$ edges, $G$ contains a cycle of length at least $n$.*

As an immediate consequence we obtain the following useful decomposition lemma of Figaj and Łuczak [7].

**Lemma 4.** *If no non-bipartite component of a graph* $G$ *contains a matching of at least* $n/2$ *edges, then there exists a partition* $V(G) = V^1 \cup V^2 \cup V^3$ *of the vertices of* $G$ *for which*

- *(A)* $G$ *has no edges joining* $V^1 \cup V^2$ *and* $V^3$*;*
- *(B) the subgraph* $G[V^1 \cup V^2]$ *is bipartite, with bipartition* $(V^1, V^2)$*;*
- *(C) the subgraph* $G[V^3]$ *has at most* $n(|V^3|-1)/2$ *edges and each component of* $G[V^3]$ *is non-bipartite.*

Notice that Lemma 4 defines a decomposition of $V(G)$ into sets $V^1$, $V^2$, and $V^3$, and we shall call $V^3$ the *sparse set*. For completeness, we include its proof.

**Proof.** Assume that no non-bipartite component of $G$ contains a matching of at least $n/2$ edges. Let $(V^1, V^2)$ be the bipartition of the union of bipartite components of $G$, and $V^3$ be the vertex set of the union of non-bipartite components. Then (A) and (B) hold, and (C) follows from Theorem 3. ∎

## 3. Odd cycles

For an integer $k \geq 1$ and a positive number $c$, let $P_k(c)$ be the following property:

> $P_k(c)$: *for every* $\varepsilon > 0$ *there exist a* $\delta > 0$ *and an* $n_0$ *such that for every odd* $n > n_0$ *and any graph* $G$ *with* $v(G) > (1+\varepsilon)cn$ *and* $e(G) \geq (1-\delta)\binom{v(G)}{2}$*, any* $k$*-edge-colouring of* $G$ *has a monochromatic non-bipartite component with a matching of* $(n+1)/2$ *edges.*

Our proof of Theorem 1 is based on the following consequence of the Regularity Lemma [14] observed by Figaj and Łuczak; see [7] for a more general statement.

**Lemma 5.** *Let a real number* $c > 0$ *and an integer* $k \geq 2$ *be given. If* $P_k(c)$ *holds, then*

$$R_k(C_n) \leq (c + o(1))n, \qquad \text{as } n \to \infty.$$

Hence, Theorem 1 follows from the next lemma.

**Lemma 6.** *For any integer $k \geq 4$, $P_k(k2^k)$ holds.*

**Proof.** Given an integer $k \geq 4$ and an $\varepsilon > 0$, let $n$ be a sufficiently large odd integer, $\delta = \varepsilon/2^{2k+4}$ and $N = (1+\varepsilon)k2^kn$. Suppose that $G$ is a graph with $v(G) \geq N$ and $e(G) \geq (1-\delta)\binom{v(G)}{2}$. Assume to the contrary that there exists a $k$-edge colouring of $G$ without a monochromatic matching of $(n+1)/2$ edges in a non-bipartite component. We may also assume that $\varepsilon < 1$ and $v(G) = N$. Indeed, if $v(G) > N$ and

$$e(G) \geq (1-\delta)\binom{v(G)}{2}, \tag{1}$$

then, iteratively removing ($v(G) - N$ times) a vertex of minimum degree, we obtain a subgraph of $G$ with $N$ vertices and at least $(1-\delta)\binom{N}{2}$ edges.

For every colour $i$, let $G_i$ be the spanning subgraph of $G$ induced by the edges coloured by $i$. Then no $G_i$ contains a matching of $(n+1)/2$ edges in a non-bipartite component, otherwise we would have a contradiction.

We apply Lemma 4 to $G_i$ for every $i \in [k] := \{1, \ldots, k\}$ and obtain a partition into $V_i^1$, $V_i^2$, and the sparse set $V_i^3$. For every $i \in [k]$, set $X_i^1 = V_i^1$ and $X_i^2 = V_i^2 \cup V_i^3$. Notice that there are $2^k$ sets of the form $\bigcap_{\ell=1}^{k} X_\ell^{j_\ell}$, where $j_\ell \in \{1,2\}$ for every $\ell$. Since $V_i^1$, $V_i^2$ and $V_i^3$ form a partition of $V(G)$ for every $i$, it is clear that the above $2^k$ sets are pairwise disjoint and form a partition of $V(G)$.

The graph $G$ has $N = (1+\varepsilon)k2^kn$ vertices, therefore, there is a choice of $j_\ell \in \{1,2\}$, $\ell = 1, 2, \ldots, k$, such that the size of the set $X = \bigcap_{\ell=1}^{k} X_\ell^{j_\ell}$ is at least $N/2^k = (1+\epsilon)kn > kn$.

For every $i$, if there is and edge $e$ of colour $i$ in $X$, then it must be contained in $V_i^3$ (by (A) and (B)). Hence, it is contained in an odd component (by (C)). Since there are no monochromatic matchings of $(n+1)/2$ edges in the non-bipartite components, $X$ contains no cycles longer than $n$ in colour $i$, so, by Theorem 3, there are at most $n(|X|-1)/2$ edges of colour $i$ with both endpoints in $X$. Hence,

$$e(G[X]) \leq kn(|X|-1)/2. \tag{2}$$

On the other hand, from (1), we have

$$e(G[X]) \geq \binom{|X|}{2} - \delta\binom{N}{2}. \tag{3}$$

Comparing (2) and (3) yields

$$|X| \leq kn + \delta\frac{N(N-1)}{|X|-1}.$$

Using assumptions $\varepsilon < 1$, $\delta = \varepsilon/2^{2k+4}$, $N \leq k2^{k+1}$, and $|X| > kn$, we have that

$$\delta\frac{N(N-1)}{|X|-1} \leq 2\delta\frac{N^2}{|X|} \leq 2\delta\frac{(k2^{k+1}n)^2}{kn} \leq \frac{\varepsilon kn}{2}.$$

Thus,

$$(1+\varepsilon)kn \leq |X| \leq kn + \frac{\varepsilon kn}{2},$$

which is a contradiction. ∎

**Remark 7.** The methods of Figaj and Łuczak and the proof above give a result slightly stronger than Theorem 1.

*Given a natural number* $k \geq 4$ *and an* $\varepsilon > 0$*, there exist a* $\delta > 0$ *and an* $n_0$ *with the following property. Suppose that* $n > n_0$ *is odd,* $N \geq (1+\varepsilon)k2^k n$*, and* $G$ *is a graph with* $v(G) \geq N$ *and* $e(G) \geq (1-\delta)\binom{v(G)}{2}$*. Then in any* $k$*-colouring of the edges of* $G$*, there exists a monochromatic cycle* $C_n$*.*

These types of theorems are not much more difficult than the ones on the colourings of the complete graphs, however, these are the forms we use in our applications.

## 4. Even cycles

In the proof of Theorem 2 we shall use another case of the lemma of Figaj and Łuczak (Lemma 3 in [7]).

**Lemma 8.** *Let a real number* $c > 0$ *be given. If for every* $\varepsilon > 0$ *there exist a* $\delta > 0$ *and an* $n_0$ *such that for every even* $n > n_0$ *and any graph* $G$ *with* $v(G) > (1+\varepsilon)cn$ *and* $e(G) \geq (1-\delta)\binom{v(G)}{2}$*, any* $k$*-edge-colouring of* $G$ *has a monochromatic component containing a matching of* $n/2$ *edges, then*

$$R_k(C_n) \leq (c + o(1))n.$$

Now we prove Theorem 2.

**Proof.** For an arbitrary $0 < \varepsilon < 1$, consider any $k$-colouring of a graph $G$ on $N > (1+\varepsilon)nk$ vertices and with at least $(1-\varepsilon/3)\binom{N}{2}$ edges. One of the colours must have at least $\frac{1}{k}(1-\varepsilon/3)\binom{N}{2} > \frac{1}{2}n(N-1)+1$ edges, so, by Theorem 3, this colour contains a cycle of length at least $n+1$. This implies the existence of a matching covering $n$ vertices in a monochromatic component. Hence, Lemma 8 implies that $R_k(C_n) \le (k+o(1))n$. ∎

# References

[1] H. Abbott and D. Hanson, *A problem of Schur and its generalizations*, Acta Arith **20** (1972), 175–187.

[2] F. Benevides and J. Skokan, *The 3-colored Ramsey number of even cycles*, J. Combin. Theory Ser. B **99** (2009), no. 4, 690–708.

[3] J. A. Bondy and P. Erdős, *Ramsey numbers for cycles in graphs*, J. Combin. Theory Ser. B **14** (1973), 46–54.

[4] F.R.K. Chung, *On triangular and cyclic Ramsey numbers with k colors*, Graphs and combinatorics (Proc. Capital Conf., George Washington Univ., Washington, D.C., 1973), pp. 236–242, Lecture Notes in Math., Vol. 406, Springer, Berlin 1974.

[5] P. Erdős and T. Gallai, *On maximal paths and circuits of graphs*, Acta Math. Acad. Sci. Hungar. **10** (1959), 337–356 (unbound insert).

[6] R. J. Faudree and R. H. Schelp, *All Ramsey numbers for cycles in graphs*, Discrete Math. **8** (1974), 313–329.

[7] A. Figaj and T. Łuczak, *The Ramsey number for a triple of large cycles*, `arXiv:0709.0048v1 [math.CO]`.

[8] Y. Kohayakawa, M. Simonovits, and J. Skokan, *The 3-colored Ramsey number of odd cycles*, Proceedings of GRACO 2005, pp. 397–402 (electronic), Electron. Notes Discrete Math., 19, Elsevier, Amsterdam, 2005.

[9] Y. Kohayakawa, M. Simonovits, and J. Skokan, *The 3-colored Ramsey number of odd cycles*, J. Combin. Theory Ser. B, to appear.

[10] Y. Li, *The multi-color Ramsey number of an odd cycle*, J. Graph Theory **62** (2009), no. 4, 324–328.

[11] Y. Li and K. Lih, *Multi-color Ramsey numbers of even cycles*, European J. Combin. **30** (2009), no. 1, 114–118.

[12] T. Łuczak, $R(C_n, C_n, C_n) \leq (4 + o(1))n$, J. Combin. Theory Ser. B **75** (1999), no. 2, 174–187.

[13] V. Rosta, *On a Ramsey-type problem of J. A. Bondy and P. Erdős. I, II*, J. Combin. Theory Ser. B **15** (1973), 94–104; ibid. 15 (1973), 105–120.

[14] E. Szemerédi, *Regular partitions of graphs*, Problèmes combinatoires et théorie des graphes (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976), Colloq. Internat. CNRS, vol. 260, CNRS, Paris, 1978, pp. 399–401.

[15] S. Yongqi, Y. Yuansheng, X. Feng, and L. Bingxi, *New lower bounds on the multicolor Ramsey numbers* $R_r(C_{2m})$, Graphs Combin. **22** (2006), no. 2, 283–288.

[16] H. Wan, *Upper bounds for Ramsey numbers* $R(3, 3, ..., 3)$ *and Schur numbers*, J. Graph Theory **26** (1997), 119–122.

[17] X. Xu, Z. Xie, G. Exoo, and S. Radziszowski, *Constructive lower bounds on classical multicolor Ramsey numbers*, Electron. J. Combin. **11** (2004), no. 1, #R35, 24pp. (electronic).